

## Management of Small Grain Diseases Fungicide Efficacy for Control of Wheat Diseases

The North Central Regional Committee on Management of Small Grain Diseases (NCERA-184) has developed the following information on fungicide efficacy for control of certain foliar diseases of wheat for use by the grain production industry in the U.S. Efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy is based on proper application timing to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are 132.32121 based on a single application of the labeled rate as listed in the table. Table includes most widely marketed products, and is not intended to be a list of all labeled products.

### Efficacy of fungicides for stripe rust control based on appropriate application timing

Fungicide(s)					
Class	Active ingredient	Product	Rate/A (fl. oz)	Stripe rust	Harvest Restriction
Strobilurin	Azoxystrobin 22.9%	Quadris 2.08 SC	6.2 - 10.8	E <sup>2</sup>	45 days
	Fluoxastrobin 40.3%	Evito 480 SC	2.0 – 4.0	-- <sup>3</sup>	40 days
	Pyraclostrobin 23.6%	Headline SC	6.0 - 9.0	E <sup>2</sup>	Feekes 10.5
Triazole	Cyproconazole 8.9%	Alto 100 SL	3.0 - 5.5	-- <sup>3</sup>	30 days
	Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	E	30 days
	Propiconazole 41.8%	Tilt 3.6 EC <sup>4</sup>	4.0	VG	Feekes 10.5
	Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	-- <sup>3</sup>	30 days
	Tebuconazole 38.7%	Folicur 3.6 F <sup>4</sup>	4.0	E	30 days
	Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.2	E	30 days
Mixed mode of action	Metconazole 7.4% Pyraclostrobin 12%	TwinLine 1.75 EC	7.0 – 9.0	E	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 7.0%	Quilt 200 SC	14.0	E	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 13.5%	Quilt Xcel 2.2 SE <sup>5</sup>	14.0	-- <sup>3</sup>	Feekes 10.5
	Propiconazole 11.4% Trifloxystrobin 11.4%	Stratego 250 EC	10.0	VG	35 days
	Tebuconazole 22.6% Trifloxystrobin 22.6%	Absolute 500 SC	5.0	-- <sup>3</sup>	35 days

<sup>1</sup> Efficacy categories: VG=Very Good; E=Excellent

<sup>2</sup> Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred

<sup>3</sup> Insufficient data to make statement about efficacy of this product

<sup>4</sup> Multiple generic products containing the active ingredients propiconazole and tebuconazole may also be labeled in some states.

Products including tebuconazole include: Embrace, Monsoon, Muscle 3.6 F, Onset, Orius 3.6 F, Tebucon 3.6 F, Tebustar 3.6 F, Tebuzol 3.6 F, Tegrol, and Toledo. However, check with local Extension personnel to determine if a specific product is labeled within the appropriate state. Products containing propiconazole include: Bumper 41.8 EC, Fitness, Propiconazole E-AG, and PropiMax 3.6 EC.

**This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions. No endorsement is intended for products listed, nor is criticism meant for products not listed. Members or participants in the NCERA-184 committee assume no liability resulting from the use of these products.**