# 2019 Mississippi On-Farm Cotton Variety Trials

## Darrin M. Dodds, Bradley Norris, Joey Williams, and William Rutland Mississippi State University Extension





### **EXTENSION**

## **Table of Contents**

2019 County Trial Locations and Cooperators	
Introduction	4
Methodology	4
Entries	5
Site Characteristics	5
Reported Data & Analysis	5
Data Tables	6
Data Summarized across All Locations	6
Data Summarized across Delta Locations	6
Data Summarized across Hill Locations	7
Data Summarized across Irrigated Locations	7
Data Summarized across Dryland Locations	8
Individual Trial Location Data:	
Brooksville	8
Bruce	9
Coffeeville	9
Crawford	
Edwards	
Ellistown	
Eupora	
Glendora	
Greenwood	
Greenwood	
Louise	
Mayersville	14
Mississippi State	14
Okolona	15
Natchez	15
Sledge	16
West Point	16

### **2019** County Trial Locations and Cooperators

Trials arranged and conducted by: Dr. Darrin Dodds

Assistance provided by: Lucas Franca, Jake McNeal, Steven Hall, Brint Lindsey, Ty Dickson, Eli Hobbs, and Wilson Whitlock

Special thanks to: Dr. Tyson Raper – University of Tennessee – West Tennessee Research and Education Center **Table 1. Locations, growers, and cooperating agronomists for 2019 Mississippi State University County** 

Variety Trial Program

Location	Grower	MSU Agronomist
Brooksville	Darrin Dodds	Darrin Dodds
Bruce	Mr. Trey Brower	Darrin Dodds
Coffeeville	Mr. Coley Bailey	Darrin Dodds
Crawford	Mr. Rodney Mast/Lowell Mullett	Darrin Dodds
Edwards	Mr. Kendall Garraway	Darrin Dodds
Ellistown	Mr. Larry Coker	Mr. Charlie Stokes
Eupora	Mr. Matt Knight	Dr. Bill Burdine
Glendora	Mr. Mike Sturdivant	Darrin Dodds
Greenwood	Mr. John Moor	Mr. Andy Braswell
Greenwood	Mr. Travis Dunn	Mr. Andy Braswell
Louise	Mr. Byron Seward	Darrin Dodds
Mayersville	Mr. Chase Mahalitic	Darrin Dodds
Mississippi State	Darrin Dodds	Darrin Dodds
Okolona	Mr. Matthew Poe	Dr. Bill Burdine
Natchez	Mr. Matthew Guedon	Darrin Dodds
Sledge	Mr. Sledge Taylor	Darrin Dodds
West Point	Mr. Ben Harlow	Mr. Charlie Stokes

Mississippi State University Extension sincerely appreciates the time and effort of the cooperating growers and Mississippi State University Agronomists. In addition, several Independent Consultants provided a tremendous level of assistance with these trials including: Mr. Ty Edwards, Mr. Jason Grafton, Mr. Bert Falkner, Mr. Tucker Miller, and Mr. Tim Richards. Sincere gratitude is also extended to the following seed companies and representatives for providing seed for these trials: BASF – Dr. Andy White, Crop Production Services/Dyna-Gro – Mr. Scott Cummings, Phytogen Cottonseed – Dr. Tom Eubank, Americot/NexGen – Mr. Chase Samples, and Delta and Pine Land – Mr. Greg Ferguson. Cooperation from all aforementioned parties is essential for success of the Mississippi State University County Research and Demonstration Yield Trial Program. In addition, partial financial support for this project was provided by each participating company and Cotton Incorporated.

### Introduction

The cotton variety selection process is often difficult and, in many cases, leaves growers wondering for the remainder of the growing season whether they made the right variety selection decisions. Furthermore, the rapid introduction of new varieties and discontinued production of "older" varieties has become commonplace over the past several years. Historically, a premier variety would remain in the marketplace for a long period of time. However, a variety that performs well today typically has a life span of four to six years. One that does not perform well will likely remain on the market for less than three years. In addition, the historical standard for variety testing information was to have two to three years of data prior to release of any given variety. Today, one to two years of "broad scale" variety testing is common prior to release of a new variety. Therefore, greater demand has been placed upon testing a variety in as many environments as possible as a substitute for multiple years of data. In most cases, variety testing prior to release is conducted by private industry through a series of testing methods and through University Official Variety Trial (OVT) programs. OVT data is typically available for one year prior to release of a given variety.

Our on-farm testing program is not designed to replace or compete with small-plot OVT testing programs, rather it is designed to complement the data that is provided by OVT programs. The use of large plot variety trial data in conjunction with small plot OVT data provides a tremendous resource with respect to variety performance to the growers of Mississippi.

### Methodology

The on-farm testing program at Mississippi State University is designed to test varieties in as many environments as possible. Limiting the number of entries allows for efficient planting and harvest operations and requires a minimum amount of time from cooperating growers. The number of variety entries each given company is given is dependent upon market share. In addition, one to two at-large entries are given to smaller companies in order to provide equal opportunity to as many seed providers as possible. Our on-farm variety tests are usually planted in 8- or 12-row sets utilizing planting equipment provided by each respective grower. In some cases, 4- or 6-row sets are used depending on site characteristics and grower preference. In addition, two replications of each variety are planted and harvested at all locations. Plot lengths ranged from 500 to 2600 feet in 2019 depending on the characteristics of the field the trial was conducted in. Seed treatments are at the discretion of the company providing seed. A premium seed treatment package including an insecticide, fungicide, and nematacide was provided for each variety. In-season management is at the discretion of the grower and each is encouraged to manage the plot area as he/she would manage any given field on their farm.

Each replication for each variety was individually harvested using standard harvest equipment. Harvest weights were collected using a boll buggy or trailer modified to display the weight of seed cotton contained therein. Prior to all harvest operations, each boll buggy or trailer was calibrated by the Mississippi Department of Agriculture to ensure that accurate harvest weights were collected. An 8- to 10-pound seed cotton sample was collected for each variety tested. In order to reduce ginning time, subsamples from replications number one and two were composited into a single sample. Seed cotton was ginned at the University of Tennessee – West Tennessee Research and Education Center. Ginning equipment at the WTREC consists of a 20-saw Continental Eagle gin equipped with a stick machine, incline cleaners, two lint cleaners, and a condenser. Fiber quality for each ginned sample was determined using a High Volume Instrument (HVI) located at the United States Department of Agriculture Classing Office in Memphis, TN.

### **Entries**

A maximum of 10 core variety entries per year are allowed in the Mississippi State University on-farm variety trial program. Entries are allotted by market share from respective companies. One entry per year is automatically given to the variety planted on the highest acreage in the previous year based on the annual Varieties Planted Report from USDA-AMS. In 2019, Monsanto/Delta and Pine Land was allotted three spots; Phytogen Cottonseed was allotted three spots, Americot was allotted two spots; and two additional "at-large" entries were given to provide parody between smaller companies with less resources than larger companies. Entries in the 2019 Mississippi State University County Trial Program were as follows:

Slot #	Criteria/Company	Variety
1	At – Large Entry – Crop Production Services/Dyna-Gro	DG 3526 B2XF
2	At – Large Entry – BASF	ST 5600B2XF
3	Delta and Pine Land	DP 1646 B2XF
4	Delta and Pine Land	DP 1835 B3XF
5	Delta and Pine Land	DP 1845 B3XF
6	Americot	NG 3994 B3XF
7	Americot	NG 4936 B3XF
8	Phytogen Cottonseed	PHY 350 W3FE
9	Phytogen Cottonseed	PHY 400 W3FE
10	Phytogen Cottonseed	PHY 480 W3FE

 Table 2. 2019 Mississippi State University County Variety Trial Program entry list.

### **Site Characteristics**

Locations for the 2019 Mississippi State University County Yield Trial Program are listed on page 3. Yield trials were conducted at a total of 17 locations. Six locations were located in the Delta and eleven were in the hills. All Delta locations were irrigated and ten of eleven Hill locations were dryland. The remaining Hill location (Crawford) was pivot irrigated. Field sites were chosen based upon grower preference and required elements to conduct a reliable yield trial.

### **Reported Data & Analysis**

Each data table includes the following: variety, lint yield, lint percent, micronaire, staple length (in inches) fiber strength, fiber uniformity, and leaf grade. Data analysis using SAS v. 9.4 was conducted on all replicated trials. Grand means (averages) are presented as well as Least Significant Differences (LSD). Least Significant Differences are the smallest value with which we can confidently say there is a difference between two means. Differences in means less than the given LSD value are likely due to variability within a given field or environment. For non-replicated trials and fiber data at individual locations, LSD's are not applicable. For locations that were replicated and data from one replication of a given variety was lost, SAS will interpret these data as missing and provide data analysis based on estimates. Therefore, average data for a given location may be slightly different than data reported.

Table 5. Yield an	a nder quan	ty data pooled	across.	I / locations.			
Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	1076*	40.2	4.4	1.18	33.0	82.2	3.5
DP 1646 B2XF	1009	39.5	4.5	1.20	31.1	82.1	3.4
DG 3526 B2XF	973	40.2	4.7	1.15	30.5	82.9	3.6
DP 1845 B3XF	968	39.2	4.2	1.23	33.4	82.6	3.8
NG 4936 B3XF	959	37.2	4.5	1.20	31.4	82.6	3.1
PHY 350 W3FE	952	37.6	4.5	1.18	32.1	82.7	3.5
ST 5600B2XF	952	39.3	4.8	1.17	32.8	82.7	3.6
DP 1835 B3XF	928	40.3	4.6	1.18	31.8	82.1	3.5
PHY 480 W3FE	904	38.2	4.5	1.18	32.3	83.1	3.5
NG 3994 B3XF	839	38.6	4.8	1.17	31.1	82.1	3.8
Grand Mean	956	39.0	4.5	1.18	31.9	82.5	3.5
LSD (0.05)	54	0.7	0.1	0.02	0.3	0.7	NS

### **Yield and Fiber Quality Data Pooled Across 17 Locations**

Table 3.	Yield and	fiber qua	ality data	pooled acr	oss 17 locations.
----------	-----------	-----------	------------	------------	-------------------

\* Yield not statistically different than the top yielding variety.

### Delta Region Locations Included: Glendora, Greenwood (2 Locations), Louise, Mayersville, and Sledge

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	10/0	
PHY 400 W3FE	1156*	39.4	4.4	1.20	34.0	82.6	3.3
DP 1646 B2XF	1050*	37.9	4.6	1.23	30.5	82.8	3.7
ST 5600B2XF	1017	39.2	4.7	1.17	33.4	82.3	3.8
NG 4936 B3XF	1007	37.3	4.5	1.21	31.5	81.8	3.0
DP 1845 B3XF	996	39.1	4.2	1.25	34.5	82.7	4.0
DG 3526 B2XF	995	39.6	4.7	1.14	30.4	82.7	3.5
PHY 350 W3FE	961	37.1	4.6	1.18	32.5	83.2	2.9
PHY 480 W3FE	959	37.9	4.4	1.19	33.1	83.2	3.3
NG 3994 B3XF	899	39.1	4.9	1.19	31.9	82.6	3.8
DP 1835 B3XF	896	39.5	4.5	1.18	32.0	82.0	3.2
Grand Mean	994	38.6	4.5	1.19	32.4	82.6	3.5
LSD (0.05)	117	1.4	0.3	0.03	1.5	NS	NS

Table 4. Yield and fiber quality data pooled over six Delta locations

#### Hill Region Locations Included: Brooksville, Bruce, Coffeeville, Crawford, Edwards, Ellistown, Eupora, Mississippi State, Okolona, Natchez, and West Point

	ind noer qua	nty data pooled		even min region	iocations.		
Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	<b>1026</b> <sup>*</sup>	40.7	4.4	1.16	32.4	82.0	3.5
DP 1646 B2XF	<b>974</b> *	40.4	4.5	1.19	31.4	81.6	3.2
DG 3526 B2XF	947	40.7	4.7	1.14	30.5	83.0	3.6
DP 1845 B3XF	939	39.4	4.3	1.21	32.7	82.5	3.6
PHY 350 W3FE	931	38.0	4.5	1.18	31.8	82.4	3.7
DP 1835 B3XF	928	40.9	4.6	1.17	31.6	82.1	3.7
NG 4936 B3XF	921	37.4	4.5	1.19	31.3	83.1	3.1
ST 5600B2XF	908	39.5	4.8	1.16	32.4	82.9	3.5
PHY 480 W3FE	866	38.5	4.5	1.17	31.9	83.0	3.5
NG 3994 B3XF	796	38.6	4.7	1.16	30.6	81.9	3.7
Grand Mean	923	39.4	4.5	1.17	31.7	82.4	3.5
LSD (0.05)	59	0.9	0.2	0.02	0.9	0.8	NS

	Table 5.	Yield and fiber	quality data	pooled over	eleven Hill	region locations.
--	----------	-----------------	--------------	-------------	-------------	-------------------

\* Yield not statistically different than the top yielding variety.

#### Irrigated Locations Included: Crawford, Glendora, Greenwood (2 Locations), Louise, Mayersville, and Sledge

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	1/0		Inches	- grams/tex -	%	
PHY 400 W3FE	<b>1137</b> <sup>*</sup>	38.6	4.2	1.21	34.0	82.6	3.3
DP 1646 B2XF	<b>1052</b> *	38.0	4.3	1.23	30.6	82.8	3.6
NG 4936 B3XF	1013	36.3	4.3	1.22	31.5	82.4	3.0
ST 5600B2XF	1010	38.1	4.5	1.18	33.2	82.5	3.9
DG 3526 B2XF	984	38.7	4.5	1.15	30.5	83.0	3.4
DP 1845 B3XF	968	38.2	3.9	1.25	34.2	82.9	3.9
PHY 480 W3FE	923	36.8	4.2	1.20	33.0	83.5	3.1
PHY 350 W3FE	918	36.2	4.4	1.18	32.4	83.4	3.1
DP 1835 B3XF	880	39.0	4.3	1.19	32.1	82.1	3.1
NG 3994 B3XF	874	38.2	4.7	1.20	31.8	82.8	3.9
Grand Mean	976	37.8	4.3	1.20	32.3	82.8	3.4
LSD (0.05)	101	1.3	0.2	0.03	1.3	NS	NS

Table 6. Yield and fiber quality data pooled over six irrigated locations

### 2019 Mississippi State University County Yield Trial Program - Flex Varieties

#### Dryland Locations Included: Brooksville, Bruce, Coffeeville, Edwards, Ellistown, Eupora, Mississippi State, Okolona, Natchez, and West Point

I WOLC I V IIII W							
Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	<b>1019</b> *	41.0	4.4	1.16	32.3	82.0	3.6
DP 1646 B2XF	<b>961</b> *	40.3	4.6	1.18	31.4	81.6	3.2
PHY 350 W3FE	947	38.3	4.5	1.18	31.8	82.3	3.7
DP 1845 B3XF	946	39.7	4.4	1.21	32.8	82.4	3.7
DG 3526 B2XF	945	41.0	4.8	1.14	30.4	82.9	3.7
DP 1835 B3XF	936	41.0	4.7	1.17	31.5	82.1	3.8
NG 4936 B3XF	906	37.8	4.5	1.18	31.3	82.9	3.1
ST 5600B2XF	897	40.0	4.8	1.16	32.4	82.9	3.4
PHY 480 W3FE	872	38.9	4.6	1.16	31.8	82.9	3.7
NG 3994 B3XF	796	38.8	4.7	1.15	30.6	81.7	3.7
Grand Mean	923	39.7	4.6	1.17	31.6	82.4	3.6
LSD (0.05)	60	0.9	0.2	0.03	1.0	0.9	NS

Table 7. Yield and fiber quality data pooled over ten dryland locations.

\* Yield not statistically different than the top yielding variety.

### **Individual Trial Location Data**

Location: Brooksville	Row width: 38"	Harvest date: November 20, 2019
Grower: Darrin Dodds	Irrigated: Dryland	Soil series: Brooksville Silty Clay
MSU Agronomist: D. Dodds	Planting date: May 22, 2019	

#### Table 8. Yield and fiber quality data at Brooksville.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
NG 4936 B3XF	<b>533</b> *	37.5	4.4	1.18	34.5	82.6	3.0
DP 1835 B3XF	<b>53</b> 1 <sup>*</sup>	39.7	4.4	1.19	33.8	82.4	3.0
PHY 400 W3FE	<b>519</b> *	38.9	4.1	1.16	33.6	82.0	5.0
DP 1845 B3XF	<b>516</b> *	38.9	4.3	1.17	33.7	81.5	3.0
DP 1646 B2XF	<b>490</b> *	38.2	4.2	1.22	34.1	81.9	4.0
PHY 350 W3FE	<b>462</b> *	35.8	4.2	1.20	34.0	83.5	4.0
DG 3526 B2XF	<b>459</b> *	39.4	4.5	1.14	32.6	83.6	6.0
ST 5600B2XF	<b>456</b> *	39.2	4.3	1.13	34.0	81.4	5.0
PHY 480 W3FE	<b>453</b> *	36.6	4.3	1.15	32.7	83.6	5.0
NG 3994 B3XF	291	38.7	4.6	1.18	32.8	83.3	5.0
Grand Mean	471	38.3	4.3	1.17	33.6	82.6	4.3
LSD (0.05)	106	•	•	•	•	•	•

Location: BruceRow width: 38"Harvest date: October 24, 2019Grower: Trey BrowerIrrigated: DrylandSoil series: Collins/Falaya SiltMSU Agronomist: D. DoddsPlanting date: May 22, 2019Loam

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	1442*	43.5	4.7	1.21	31.8	81.4	4.0
DG 3526 B2XF	1392*	45.4	4.4	1.19	32.0	82.8	4.0
PHY 350 W3FE	1338*	40.9	4.2	1.17	32.1	83.1	4.0
DP 1646 B2XF	1298*	43.8	4.5	1.12	31.6	81.6	4.0
ST 5600B2XF	1273	41.7	5.1	1.09	28.2	82.3	3.0
DP 1835 B3XF	1220	41.9	4.7	1.17	33.0	82.2	5.0
DP 1845 B3XF	1220	40.6	4.7	1.18	30.5	82.4	2.0
NG 3994 B3XF	1216	38.8	4.8	1.14	30.9	82.5	3.0
PHY 480 W3FE	1142	41.4	5.1	1.14	31.2	80.7	3.0
NG 4936 B3XF	829	32.0	4.4	1.16	31.6	83.2	5.0
Grand Mean	1237	41.0	4.7	1.16	31.3	82.2	3.7
LSD (0.05)	150	•	•	•	•	•	•

#### Table 9. Yield and fiber quality data at Bruce.

\* Yield not statistically different than the top yielding variety.

Location: Coffeeville	Row width: 38"	Harvest date: October 29, 2019
Grower: Coley Bailey	Irrigated: Dryland	Soil series: Collins Silt Loam
MSU Agronomist: D. Dodds	Planting date: May 24, 2019	

#### Table 10. Yield and fiber quality data at Coffeeville.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	- Lbs/Acre -	%		Inches	- grams/tex -	%	
PHY 400 W3FE	1454	39.6	4.1	1.21	34.2	83.5	4.0
NG 4936 B3XF	1370	35.5	4.1	1.22	31.7	83.8	4.0
DP 1646 B2XF	1287	39.3	4.2	1.24	32.1	80.4	3.0
PHY 350 W3FE	1285	37.6	4.4	1.18	32.3	83.5	4.0
NG 3994 B3XF	1267	38.3	5.0	1.20	32.1	83.0	5.0
DP 1845 B3XF	1239	37.4	3.9	1.27	33.9	82.2	5.0
DG 3526 B2XF	1237	40.1	4.8	1.12	28.3	82.7	3.0
DP 1835 B3XF	1192	40.1	4.6	1.19	33.2	83.0	3.0
PHY 480 W3FE	1132	39.0	3.9	1.18	32.1	82.5	5.0
ST 5600B2XF	1131	37.9	4.9	1.20	33.9	83.6	4.0
Grand Mean	1259	38.5	4.4	1.20	32.4	82.8	4.0
LSD (0.05)	NS	•	•	•	•	•	•

Location: Crawford	MSU Agronomist: D. Dodds	Planting date: May 23, 2019
Grower: Rodney Mast/Lowell	Row width: 30"	Harvest date: November 13, 2019
Mullett	Irrigated: Pivot	Soil series: Vaiden Silty Clay

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1646 B2XF	<b>1056</b> *	41.4	4.0	1.24	31.3	81.7	3.0
PHY 400 W3FE	<b>1033</b> *	37.6	3.7	1.21	33.8	81.8	3.0
NG 4936 B3XF	<b>1020</b> <sup>*</sup>	33.7	3.9	1.26	31.4	85.1	3.0
ST 5600B2XF	960 <sup>*</sup>	35.0	4.2	1.20	32.3	82.9	4.0
DG 3526 B2XF	<b>910</b> *	37.2	4.1	1.20	31.5	83.9	3.0
DP 1845 B3XF	811	36.4	3.5	1.24	32.5	83.1	3.0
DP 1835 B3XF	782	39.1	4.1	1.21	32.9	82.5	3.0
PHY 480 W3FE	750	34.2	4.0	1.20	32.5	84.3	2.0
NG 3994 B3XF	736	36.5	4.6	1.26	31.3	83.4	4.0
PHY 350 W3FE	705	34.4	4.1	1.20	31.8	83.3	4.0
Grand Mean	876	36.5	4.0	1.22	32.1	83.2	3.2
LSD (0.05)	164	•	•	•	•	•	•

#### Table 11. Yield and fiber quality data at Crawford.

\* Yield not statistically different than the top yielding variety.

Location: Edwards	Row width: 38"	Harvest date: October 29, 2019
Grower: Kendall Garraway	Irrigated: Dryland	Soil series: Reidtown Silt Loam
MSU Agronomist: D. Dodds	Planting date: May 24, 2019	

#### Table 12. Yield and fiber quality data at Edwards.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	<b>950</b> *	43.3	4.5	1.16	32.0	81.5	3.0
DG 3526 B2XF	862	39.9	4.6	1.12	30.7	83.4	3.0
DP 1835 B3XF	857	41.3	4.7	1.15	29.3	80.2	4.0
DP 1646 B2XF	841	39.7	4.6	1.19	30.2	82.9	4.0
PHY 350 W3FE	814	38.1	4.5	1.14	31.4	81.1	4.0
NG 4936 B3XF	810	36.8	4.5	1.19	30.9	81.8	3.0
ST 5600B2XF	806	37.4	4.8	1.19	33.5	82.9	4.0
PHY 480 W3FE	781	37.7	4.4	1.16	32.6	83.3	4.0
DP 1845 B3XF	771	38.1	4.0	1.23	32.0	83.4	4.0
NG 3994 B3XF	714	38.7	4.7	1.12	29.4	80.8	3.0
Grand Mean	820	39.1	4.5	1.17	31.2	82.1	3.6
LSD (0.05)	80	•	•	•	•	•	•

Location: Ellistown	Row width: 38"	Harvest date: October 24, 2019
Grower: Larry Coker	Irrigated: Dryland	Soil series: Mantachie/Talla Silt
MSU Agronomist: C. Stokes	Planting date: May 22, 2019	Loam

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	1365*	43.9	4.6	1.14	33.2	83.8	4.0
DP 1646 B2XF	1322*	42.9	4.8	1.21	30.2	81.8	3.0
DP 1835 B3XF	1308*	42.4	4.9	1.15	31.8	83.0	5.0
NG 4936 B3XF	<b>1288</b> <sup>*</sup>	39.6	4.7	1.19	31.4	84.4	4.0
DP 1845 B3XF	<b>1250</b> *	43.1	4.6	1.20	33.4	83.9	5.0
DG 3526 B2XF	1178	43.3	5.2	1.11	30.6	82.8	4.0
ST 5600B2XF	1160	40.7	5.3	1.20	33.1	82.5	3.0
PHY 350 W3FE	1147	40.0	5.0	1.18	32.4	83.1	3.0
NG 3994 B3XF	1111	42.4	5.2	1.13	30.7	80.1	3.0
PHY 480 W3FE	1092	40.2	4.9	1.16	31.8	84.7	4.0
Grand Mean	1222	41.8	4.9	1.17	31.9	83.0	3.8
LSD (0.05)	154	•	•	•	•	•	•

#### Table 13. Yield and fiber quality data at Ellistown.

\* Yield not statistically different than the top yielding variety.

Location: Eupora	Row width: 38"	Harvest date: November 14, 2019
Grower: Matt Knight	Irrigated: Dryland	Soil series: Oaklimter Silt Loam
MSU Agronomist: B. Burdine	Planting date: May 23, 2019	

#### Table 14. Yield and fiber quality data at Eupora.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
NG 4936 B3XF	1154	38.1	4.5	1.21	29.8	83.9	3.0
DP 1646 B2XF	1118	39.2	4.5	1.17	29.8	80.9	4.0
ST 5600B2XF	1113	41.1	5.0	1.18	31.4	84.5	3.0
PHY 400 W3FE	1066	41.6	4.3	1.17	31.3	81.8	3.0
PHY 350 W3FE	1057	38.4	4.5	1.18	31.4	82.7	5.0
DP 1835 B3XF	1026	41.0	4.5	1.16	29.4	81.4	4.0
PHY 480 W3FE	1009	39.5	4.5	1.15	30.7	83.9	4.0
DG 3526 B2XF	972	39.9	4.8	1.13	28.5	84.0	4.0
NG 3994 B3XF	930	39.6	4.9	1.15	28.5	80.9	4.0
DP 1845 B3XF	909	38.6	4.2	1.25	31.5	81.9	4.0
Grand Mean	1035	39.7	4.6	1.18	30.2	82.6	3.8

Location: Glendora Grower: Mike Sturdivant Jr. MSU Agronomist: D. Dodds Row width: 38" Irrigated: Furrow Planting date: May 28, 2019 Harvest date: November 5, 2019 Soil series: Dundee Silt Loam

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre-	%		Inches	- grams/tex -	%	
PHY 400 W3FE	910	40.0	4.1	1.19	34.5	83.3	4.0
DP 1646 B2XF	907	36.5	4.2	1.26	31.2	83.3	5.0
NG 4936 B3XF	873	34.7	4.1	1.22	28.2	82.5	4.0
ST 5600B2XF	831	36.1	4.3	1.23	34.9	81.5	4.0
PHY 350 W3FE	825	36.7	4.1	1.17	31.9	82.5	4.0
DG 3526 B2XF	780	38.0	4.3	1.16	31.3	83.0	4.0
PHY 480 W3FE	727	34.1	3.9	1.23	34.5	82.4	4.0
DP 1835 B3XF	724	38.5	4.3	1.22	31.7	80.6	3.0
DP 1845 B3XF	720	36.4	3.7	1.26	33.1	82.7	4.0
NG 3994 B3XF	701	34.8	4.5	1.21	31.2	82.5	4.0
Grand Mean	800	36.6	4.2	1.22	32.3	82.4	4.0

Table 15.	Yield and	fiber	quality	data	at	Glendora.
I abic 15.	i iciu anu	IIDUI	quanty	uata	aı	Official de la construction de l

Location: Greenwood	Row width: 38"	Har
Grower: John Moor	Irrigated: Furrow	Soil
MSU Agronomist: A. Braswell	Planting date: May 25, 2019	;

#### Harvest date: October 12, 2019 Soil series: Dubbs Loam/Tensas Silty Clay Loam

#### Table 16. Yield and fiber quality data at Greenwood.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	<b>1405</b> *	38.8	4.9	1.23	35.6	81.9	1.0
ST 5600B2XF	1227	41.2	4.1	1.10	32.4	80.1	4.0
PHY 350 W3FE	1201	37.6	4.7	1.17	31.3	82.7	2.0
DG 3526 B2XF	1117	40.2	4.9	1.18	31.1	82.9	3.0
PHY 480 W3FE	1102	38.0	4.6	1.23	33.7	85.2	2.0
DP 1845 B3XF	1096	40.2	4.5	1.30	33.8	84.2	3.0
NG 4936 B3XF	1078	37.0	4.9	1.24	31.4	82.3	1.0
DP 1646 B2XF	990	34.0	4.8	1.24	30.8	83.5	3.0
DP 1835 B3XF	961	40.8	4.2	1.21	33.9	83.3	3.0
NG 3994 B3XF	851	37.6	5.0	1.24	33.4	82.2	3.0
Grand Mean	1103	38.6	4.7	1.21	32.7	82.8	2.5
LSD (0.05)	110	•	•	•	•	•	•

Location: GreenwoodRow width: 38"Harvest date: October 17, 2019Grower: Travis DunnIrrigated: IrrigatedSoil series: Dundee Loam/MSU Agronomist: A. BraswellPlanting date: May 25, 2019Tensas Silty Clay Loam

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1646 B2XF	1202	39.6	4.3	1.21	30.7	82.3	4.0
DG 3526 B2XF	1157	39.5	4.3	1.13	30.5	80.8	3.0
DP 1845 B3XF	1148	38.1	3.8	1.23	35.4	79.6	5.0
NG 4936 B3XF	982	38.2	4.4	1.21	34.0	82.6	4.0
ST 5600B2XF	899	38.0	4.4	1.23	33.3	83.3	5.0
NG 3994 B3XF	898	39.5	4.6	1.18	31.6	81.8	5.0
DP 1835 B3XF	727	34.1	4.3	1.20	30.9	81.2	4.0
Grand Mean	933	37.4	4.3	1.21	32.2	82.2	4.1
LSD (0.05)	NS	•	•	•	•	•	•

#### Table 17. Yield and fiber quality data at Greenwood.

Location: Louise	Row width: 30" 2x1 Skip	Harvest date: October 14, 2019
Grower: Byron Seward	Irrigated: Furrow	Soil series: Forestadle-Brittain
MSU Agronomist: D. Dodds	Planting date: May 23, 2019	Silt Loam

#### Table 18. Yield and fiber quality data at Louise.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1646 B2XF	<b>1652</b> *	38.5	4.5	1.26	31.5	82.0	4.0
PHY 400 W3FE	<b>1626</b> <sup>*</sup>	39.8	4.3	1.20	33.8	83.3	3.0
NG 3994 B3XF	<b>1595</b> *	40.6	5.0	1.21	33.5	84.0	3.0
DP 1835 B3XF	1523	40.6	4.7	1.17	32.6	83.0	3.0
NG 4936 B3XF	1516	37.7	4.3	1.23	32.6	81.4	3.0
ST 5600B2XF	1515	39.0	5.1	1.17	33.7	83.4	3.0
DP 1845 B3XF	1505	39.7	4.0	1.25	34.3	83.9	4.0
DG 3526 B2XF	1451	40.9	4.7	1.12	29.4	83.3	3.0
PHY 480 W3FE	1445	38.7	4.5	1.17	31.9	82.0	2.0
PHY 350 W3FE	1371	36.9	4.8	1.20	32.7	84.3	3.0
Grand Mean	1520	39.2	4.6	1.20	32.6	83.1	3.1
LSD (0.05)	111	•	•	•	•	•	•

Location: Mayersville	Row width: 38"	Harvest date: November 20, 2019
Grower: Chase Mahalitic	Irrigated: Furrow	Soil series: Commerce Silty Clay
MSU Agronomist: D. Dodds	Planting date: May 29, 2019	Loam/Tunica Clay

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
PHY 400 W3FE	639	39.9	4.5	1.21	34.2	83.6	4.0
PHY 480 W3FE	625	40.1	4.3	1.20	34.6	83.6	4.0
NG 4936 B3XF	620	37.7	4.5	1.19	31.8	80.3	3.0
DP 1835 B3XF	578	41.5	4.5	1.15	32.1	81.9	3.0
ST 5600B2XF	550	39.4	5.2	1.14	34.2	82.1	4.0
PHY 350 W3FE	547	37.8	4.7	1.19	34.7	84.2	3.0
DP 1845 B3XF	531	39.3	4.4	1.25	36.0	83.6	4.0
DP 1646 B2XF	529	39.6	4.7	1.21	30.2	83.3	3.0
DG 3526 B2XF	490	40.2	5.1	1.13	30.5	83.1	5.0
NG 3994 B3XF	443	40.7	4.9	1.17	29.8	82.2	3.0
Grand Mean	555	39.6	4.7	1.18	32.8	82.8	3.6

Location: Mississippi State	Row width: 38"	Harvest date: November 18, 2019
Grower: Darrin Dodds	Irrigated: Dryland	Soil series: Catalpa/Leeper Silty
MSU Agronomist: D. Dodds	Planting date: May 20, 2019	Clay Loam

### Table 20. Yield and fiber quality data at Mississippi State.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1646 B2XF	562	39.4	4.3	1.20	32.1	80.8	3.0
PHY 400 W3FE	551	39.8	4.3	1.12	31.0	80.4	3.0
DG 3526 B2XF	532	41.6	4.6	1.13	30.6	80.2	3.0
NG 4936 B3XF	524	39.0	4.4	1.18	31.2	82.6	3.0
PHY 350 W3FE	513	37.2	4.3	1.15	31.9	81.5	2.0
PHY 480 W3FE	442	38.0	4.5	1.17	32.3	83.0	3.0
ST 5600B2XF	431	40.4	4.7	1.15	33.4	83.4	3.0
DP 1845 B3XF	421	38.7	4.1	1.20	33.9	81.1	3.0
DP 1835 B3XF	411	40.9	4.6	1.14	31.1	81.9	3.0
NG 3994 B3XF	295	39.7	4.3	1.12	30.3	81.1	3.0
Grand Mean	468	39.5	4.4	1.16	31.8	81.6	2.9
LSD (0.05)	NS	•	•	•	•	•	•

Location: Okolona	Row width: 38"	Harvest date: November 6, 2019
Grower: Matthew Poe	Irrigated: Dryland	Soil series: Leeper Silty Clay
MSU Agronomist: B. Burdine	Planting date: May 23, 2019	Loam

\_\_\_\_\_

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1835 B3XF	<b>1081</b> *	44.6	4.9	1.16	31.6	81.3	4.0
PHY 350 W3FE	<b>1045</b> *	41.0	5.2	1.15	32.6	82.6	4.0
PHY 400 W3FE	<b>1044</b> *	41.5	4.5	1.14	32.9	80.9	3.0
DP 1646 B2XF	1011	42.7	5.0	1.12	29.5	80.4	2.0
PHY 480 W3FE	977	39.8	4.5	1.13	31.8	82.3	3.0
DP 1845 B3XF	971	39.7	4.4	1.21	33.9	82.9	4.0
DG 3526 B2XF	931	42.2	5.1	1.10	30.4	82.9	4.0
ST 5600B2XF	902	39.1	4.8	1.16	32.5	83.1	3.0
NG 4936 B3XF	831	39.7	4.6	1.15	30.4	81.8	2.0
NG 3994 B3XF	794	40.2	4.7	1.11	30.0	81.1	4.0
Grand Mean	959	41.1	4.8	1.14	31.6	81.9	3.3
LSD (0.05)	70	•	•	•	•	•	

#### Table 21. Yield and fiber quality data at Okolona.

\* Yield not statistically different than the top yielding variety.

Location: Natchez	Row width: 38"	
Grower: Matthew Guedon	Irrigated: Dryland	Harvest date: October 3, 2019
MSU Agronomist: D. Dodds	Planting date: May 2, 2019	Soil series: Convent Silt Loam

#### Table 22. Yield and fiber quality data at Natchez.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1845 B3XF	996	38.3	4.4	1.22	33.2	83.1	4.0
DG 3526 B2XF	977	36.3	4.6	1.24	31.5	84.1	3.0
PHY 350 W3FE	853	36.2	4.1	1.28	31.3	80.0	4.0
ST 5600B2XF	840	40.1	4.5	1.21	32.4	82.5	3.0
PHY 400 W3FE	826	37.9	4.5	1.11	32.0	81.7	4.0
NG 4936 B3XF	826	37.5	4.6	1.17	31.3	81.7	2.0
DP 1835 B3XF	804	36.2	4.5	1.18	32.4	84.3	4.0
DP 1646 B2XF	796	36.8	4.6	1.18	34.1	82.6	3.0
PHY 480 W3FE	755	36.7	4.5	1.23	31.1	82.3	4.0
NG 3994 B3XF	687	36.2	4.2	1.22	32.9	83.0	4.0
Grand Mean	836	37.2	4.5	1.20	32.2	82.5	3.5
LSD (0.05)	NS	•	•	•	•	•	•

Location: Sledge	Row width: 38"	Harvest date: November 18, 2019
Grower: Sledge Taylor	Irrigated: Irrigated	Soil series: Falaya/Collins Silt
MSU Agronomist: D. Dodds	Planting date: May 24, 2019	Loam

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1646 B2XF	$1080^*$	41.8	4.8	1.19	28.3	82.6	3.0
ST 5600B2XF	<b>1032</b> *	39.5	5.3	1.16	31.6	83.3	3.0
NG 4936 B3XF	<b>1032</b> *	38.0	4.8	1.17	30.8	81.8	3.0
PHY 400 W3FE	<b>1031</b> *	40.2	4.6	1.18	31.7	81.0	4.0
DP 1835 B3XF	949 <sup>*</sup>	41.9	5.0	1.13	30.9	81.7	3.0
NG 3994 B3XF	945 <sup>*</sup>	41.2	5.1	1.14	31.7	82.9	5.0
DP 1845 B3XF	871	39.4	4.6	1.20	34.6	82.4	4.0
DG 3526 B2XF	870	38.2	4.9	1.12	29.3	82.9	3.0
PHY 480 W3FE	860	38.5	5.0	1.13	30.7	82.9	4.0
PHY 350 W3FE	803	37.3	4.9	1.16	31.9	82.6	2.0
Grand Mean	947	39.6	4.9	1.16	31.2	82.4	3.4
LSD (0.05)	160						

Table 23. Yield and fiber quality data at Sledge.

\* Yield not statistically different than the top yielding variety.

Location: West Point
Grower: Ben Harlow
MSU Agronomist: C. Stokes

Row width: 30" Irrigated: Dryland Planting date: May 21, 2019 Harvest date: October 11, 2019 Soil series: Houston Clay

#### Table 24. Yield and fiber quality data at West Point.

Variety	Lint Yield	Lint Percent	Mic	Staple	Strength	Uniformity	Leaf
	Lbs/Acre	%		Inches	- grams/tex -	%	
DP 1845 B3XF	993 <sup>*</sup>	42.9	4.9	1.15	31.5	81.5	3.0
PHY 400 W3FE	<b>849</b> *	41.2	4.8	1.13	30.9	83.1	3.0
PHY 350 W3FE	<b>842</b> <sup>*</sup>	39.2	5.0	1.14	28.8	82.3	3.0
PHY 480 W3FE	<b>833</b> *	41.7	4.9	1.15	32.0	82.2	2.0
DP 1835 B3XF	811	42.2	4.9	1.16	29.3	80.9	3.0
ST 5600B2XF	789	42.3	5.0	1.10	31.6	82.5	3.0
DP 1646 B2XF	764	40.9	4.8	1.17	30.5	82.9	2.0
NG 4936 B3XF	762	40.6	4.9	1.16	29.9	83.0	2.0
DG 3526 B2XF	738	41.3	5.1	1.11	28.9	82.9	3.0
NG 3994 B3XF	593	34.4	5.0	1.09	28.0	81.5	3.0
Grand Mean	797	40.7	4.9	1.14	30.1	82.3	2.7
LSD (0.05)	167	•	•	•	•	•	•