

Pre-harvest interval (restriction) of fungicides commonly used in our production system

Fungicide(s)				Harvest Restriction
Class	Active ingredient	Product	Rate/A (fl. oz)	
Strobilurin	Azoxystrobin 22.9%	Quadris 2.08 SC	6.2 - 10.8	45 days
	Fluoxastrobin 40.3%	Evito 480 SC	2.0 – 4.0	40 days
	Pyraclostrobin 23.6%	Headline SC	6.0 - 9.0	Feekes 10.5
Triazole	Cyproconazole 8.9%	Alto 100 SL	3.0 - 5.5	30 days
	Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	30 days
	Propiconazole 41.8%	Tilt 3.6 EC ⁴	4.0	Feekes 10.5
	Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	30 days
	Tebuconazole 38.7%	Folicur 3.6 F ⁴	4.0	30 days
	Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.2	30 days
Mixed mode of action	Metconazole 7.4% Pyraclostrobin 12%	TwinLine 1.75 EC	7.0 – 9.0	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 7.0%	Quilt 200 SC	14.0	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 13.5%	Quilt Xcel 2.2 SE ⁵	14.0	Feekes 10.5
	Propiconazole 11.4% Trifloxystrobin 11.4%	Stratego 250 EC	10.0	35 days
	Tebuconazole 22.6% Trifloxystrobin 22.6%	Absolute 500 SC	5.0	35 days