

Table 3. Soybean fungicide trial

Treatment ^z , rate (fl oz/A) @ Soybean stage	Severity rating (0-9)			Phytotoxicity rating (0-100%)			Cercospora blight		% Green Stem	Yield (bu/A) ^x
	Frogeye leaf spot						17	14	25	25
	1 Aug ^y	17 Aug	14 Sep	1 Aug	17 Aug	14 Sep	Aug	Sep	Sep	Sep
Nontreated	2.3 a	7.0	8.3	0.0	0.0	0.0	5.0	7.0	1.0 d	54.6 c
Copper hydroxide 2 lb @ R3	1.5 bc	6.0	8.0	0.0	0.0	0.0	5.0	6.8	3.5 cd	56.7 c
Domark 230 ME 4 oz @ R3	1.0 c	6.3	7.8	0.0	0.0	0.0	5.0	6.5	16.8 bc	61.9 ab
Mancozeb 2 lb @ R3	1.3 bc	6.5	8.5	6.3	7.5	7.5	5.0	7.0	11.0 bcd	58.3 bc
Quadris 2.08 SC 4 oz + Dimilin 2L 2 oz @ R3	1.8 ab	6.5	8.3	0.0	0.0	1.0	5.0	7.0	5.5 cd	58.2 bc
Quadris 2.08 SC 4 oz + Manniplus B Moly 2 qt @ R3	1.8 ab	7.0	8.5	8.0	0.0	0.0	5.0	7.0	1.8 cd	53.9 c
Quadris 2.08 SC 4 oz + Tilt 3.6 EC 4 oz + Topsin 4.5 FL 10 oz @ R3	1.3 bc	6.0	8.3	2.0	5.0	1.0	5.0	6.5	22.5 b	65.4 a
Stratego YLD 4.18 SC 4 oz + Belt SC 2 oz @ R3	1.0 c	6.0	8.5	19.8	13.8	6.3	4.8	6.8	8.5 bcd	58.6 bc
Viathon 4.1 SC 2 pt @ R3	1.3 bc	6.5	8.0	28.8	23.8	11.3	5.0	6.3	38.8 a	58.8 bc
LSD (0.05)	0.7	0.7	1.3	21.8	23.6	14.5	0.2	0.6	15.5	5.1
CV (%)	31.1	8.0	11.1	207.3	291.0	330.3	3.4	6.0	87.3	6.0
<i>P</i> -value for F-statistic	0.0103	0.0489	0.9444	0.1020	0.4191	0.6663	0.4613	0.1068	0.0006	0.0038

^z All fungicide treatments included a non-ionic surfactant at 0.25% v/v.

^y Means followed by the same letter(s) within a column are not significantly different according to Fisher's Protected LSD ($P=0.05$).

^x Yields are weight of soybean with moisture content adjusted to 13%.

NOTE: highlighted product or product combinations include products that are **NOT** currently labeled for application to soybean.