RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.





Section 18 Emergency Exemption

TO CONTROL RICE STINK BUG IN RICE IN THE STATE OF MISSISSIPPI

EFFECTIVE DATE: July 14, 2023 EXPIRATION DATE: October 15, 2023

Endigo® ZCX EPA Reg. No. 100-1458

EPA File Number: 23MS01

KEEP OUT OF REACH OF CHILDREN.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Endigo ZCX label (EPA Reg. No. 100-1458).

This label must be in the possession of the user at the time of application.

Any adverse effects resulting from the use of Endigo ZCX under this exemption must be immediately reported to the Mississippi Department of Agriculture.

CROP USE DIRECTIONS

Crop	Pest	Rate of Endigo ZCX per Acre per Application	Rate of Active Ingredient per Acre per Application
Rice	Rice Stink Bug	4.5 fl oz	0.063 lb ai thiamethoxam + 0.03 lb ai lambda-cyhalothrin
		to	to
		6.0 fl oz	0.084 lb ai thiamethoxam + 0.04 lb ai lambda-cyhalothrin

Use Restrictions:

- To reduce risks to bees due to spray drift, the area within 100 feet of the edge of the field (paddy) must be kept free of flowering weeds.
- **Application Timing:** Apply as required by scouting. Base timing of applications upon insect populations reaching locally determined economic thresholds.
- Application Rate: One application may be made per year at a rate of 4.5 to 6.0 fl oz per acre.
- **Maximum Allowed:** Do not exceed a total of 6.0 fl. oz. of Endigo ZCX (0.084 lb of thiamethoxam and 0.04 lb of lambda-cyhalothrin) per acre per year. Do not exceed a total of 0.12 lb a.i. of **any** lambda-cyhalothrin-containing products per acre per year.
- Pre-Harvest Interval (PHI): 21 days
- Water Volume: Apply by ground or air equipment using sufficient water volume to ensure thorough coverage of foliage. Do not use less than 2 GPA for aerial applications. Do not apply as an ultra-low volume (ULV) spray. Adding an emulsified crop oil (e.g. 1 pt. per acre) when lower aerial application volumes are used is recommended to help improve coverage, reduce evaporation, and improve efficacy.
- Flood Water: Do not release flood water within 7 days of an application.
- Aquaculture: Do not use treated rice fields for the aquaculture of edible fish and crustacea.

©2023 Syngenta

Endigo® and the Syngenta logo are trademarks of a Syngenta Group Company

Registrant: Syngenta Crop Protection, LLC PO Box 18300 Greensboro, NC 27419-8300

Label Code: MS1458052AB0523